

University of Dundee

Community-acquired pneumonia related to intracellular pathogens

Cillóniz, Catia; Torres, Antoni; Niederman, Michael; van der Eerden, Menno; Chalmers, James; Welte, Tobias

Published in:
Intensive Care Medicine

DOI:
[10.1007/s00134-016-4394-4](https://doi.org/10.1007/s00134-016-4394-4)

Publication date:
2016

Document Version
Peer reviewed version

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Cillóniz, C., Torres, A., Niederman, M., van der Eerden, M., Chalmers, J., Welte, T., & Blasi, F. (2016). Community-acquired pneumonia related to intracellular pathogens. *Intensive Care Medicine*, 42(9), 1374-1386. <https://doi.org/10.1007/s00134-016-4394-4>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Figure 1. Distribution of microbial etiology on CAP patients: Ref. [22]




Figure 2. Notification rates of Legionnaires’ disease per million by sex and age group, EU/EEA, 2014




Figure 3. Reported cases and notifications of Legionnaires’ disease per million, by reporting country, EU/EEA, 2014






Figure 4. Age and sex distribution of patients with *Mycoplasma pneumoniae*



Age and sex distribution of patients with *Mycoplasma pneumoniae*–positive respiratory tract samples (n = 96), Germany, March 2011–December 2012. Percentage of positive samples for each age group: 18–29 y, 28.1%; 30–39 y, 21.9%; 40–49 y, 22.9%; 50–59 y, 14.6%; >60 y, 13.5%. (Total >100% due to rounding.)
Ref. Dumke et al.[8].

Figure 5. Microbial Diagnosis of Atypical pathogens



Microorganism, Specimen types, diagnostic tests